






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : A61K 37/02, C07K 5/06, 5/08 C07K 5/10, C07D 405/00 C07D 207/00		A1	(11) International Publication Number: WO 89/ 03223 (43) International Publication Date: 20 April 1989 (20.04.89)
(21) International Application Number: PCT/US88/03506 (22) International Filing Date: 6 October 1988 (06.10.88) (31) Priority Application Number: 105,768 (32) Priority Date: 6 October 1987 (06.10.87) (33) Priority Country: US		(81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), DK, FI, FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), NO, SE (European patent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71)(72) Applicants and Inventors: BACHOVCHIN, William, W. [US/US]; 71 Warwick Road, Melrose, MA 02176 (US). PLAUT, Andrew, G. [US/US]; 22 Peacock Farm Road, Lexington, MA 02173 (US). KETTER, Charles, A. [US/US]; 2411 Chatham Drive, Wilmington, DE 19803 (US). (74) Agent: CLARK, Paul, T.; Fish & Richardson, One Financial Center, Suite 2500, Boston, MA 02111-2658 (US).			





(54) Title: PROTEASE INHIBITORS




Propyl Boronate



Prolyl phosphonate



Prolyl Trifluoro alkyl ketone



(57) Abstract

A compound having the structure (I), where T is of formula (II), where each D¹ and D², independently, is a hydroxyl group, is capable of being hydrolyzed to a hydroxyl group in aqueous solution at physiological pH; a

arising about a carbonyl arene and a phosphorus heteroatom. X is a group having at least one amino acid, Y is (VI) or (VII) or (VIII) and each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 is separately a group which does not interfere significantly (i.e., does not lower the K_i of the compound to less than $10^{-7}M$) with site-specific recognition of the compound by the enzyme, and allows a complex to be formed with the enzyme.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CA Canada	CZ Czech Republic	
CH Switzerland	DE Germany, Federal Republic of	
CL Chile	DK Denmark	
CO Colombia	EE Estonia	
DE Germany, Federal Republic of	FI Finland	
DK Denmark	FR France	
FI Finland	GB United Kingdom	
	GR Greece	
	GU Guernsey	
	HK Hong Kong	
	IL Israel	
	IN India	
	IR Iran	
	IS Iceland	
	IT Italy	
	JE Jersey	
	JO Jordan	
	KE Kenya	
	KH Cambodia	
	KR South Korea	
	KG Kyrgyzstan	
	KI Kiribati	
	KM Comoros	
	KN St. Kitts and Nevis	
	KO North Korea	
	KR South Korea	
	KU Kuwait	
	KV Kazakhstan	
	KY Cayman Islands	
	KZ Kazakhstan	
	LA Laos	
	LB Lebanon	
	LC St. Lucia	
	LI Liechtenstein	
	LK Sri Lanka	
	LR Liberia	
	LS Lesotho	
	LT Lithuania	
	LU Luxembourg	
	LV Latvia	
	LY Libya	
	MA Morocco	
	MC Monaco	
	MD Moldova	
	ME Montenegro	
	MG Madagascar	
	MH Marshall Islands	
	MI Micronesia	
	ML Mali	
	MM Myanmar	
	MN Mongolia	
	MO Macao	
	MP Northern Mariana Islands	
	MQ Martinique	
	MR Mauritania	
	MS Montserrat	
	MU Mauritius	
	MV Maldives	
	MW Malawi	
	MX Mexico	
	MY Malaysia	
	MZ Mozambique	
	NA Namibia	
	NC New Caledonia	
	NE Niger	
	NG Nigeria	
	NL Netherlands	
	NI Nicaragua	
	NO Norway	
	NP Nepal	
	NR Nauru	
	NU Niue	
	NZ New Zealand	
	OA Oman	
	OC Oceania	
	OD Outer Mongolia	
	OE European Union	
	OF Overseas Countries and Territories	
	OG Omani Gulf	
	OH Ohio	
	OJ Oregon	
	OK Oklahoma	
	OL Ontario	
	OM Oman	
	ON Ontario	
	OO Other	
	OP Overseas Possessions	
	OS Oregon	
	OT Overseas Territory	
	OU Overseas Union	
	OV Overseas Village	
	OW Overseas Work	
	OX Overseas X-ray	
	OY Overseas Yacht	
	Pakistan	
	Panama	
	Papua New Guinea	
	Paraguay	
	Peru	
	Philippines	
	Puerto Rico	
	Romania	
	Russia	
	Saudi Arabia	
	Senegal	
	Seychelles	
	Sierra Leone	
	Singapore	
	Slovakia	
	Slovenia	
	South Africa	
	Spain	
	St. Helena	
	St. James	
	St. John	
	St. Peter	
	St. Vincent	
	St. William	
	Sweden	
	Switzerland	
	Taiwan	
	Tanzania	
	Togo	
	Tonga	
	Turkey	
	Turkmenistan	
	USA	
	United States of America	
	Uruguay	
	Uzbekistan	
	Venezuela	
	Zambia	
	Zimbabwe	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CA Canada	LA Laos	
CH Switzerland	LV Latvia	
CN China	LT Lithuania	
DE Germany, Federal Republic of	LU Luxembourg	
DK Denmark	MC Monaco	US United States of America
FI Finland	MG Madagascar	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CA Canada	CZ Czech Republic	
CH Switzerland	DE Germany, Federal Republic of	
CL Chile	DK Denmark	
CO Colombia	EE Estonia	
DE Germany, Federal Republic of	FI Finland	
DK Denmark	FR France	
FI Finland	GB United Kingdom	
	GR Greece	
	GU Guadeloupe	
	HK Hong Kong	
	IL Israel	
	IN India	
	IR Iran	
	IS Iceland	
	IT Italy	
	JP Japan	
	KE Kenya	
	KR Korea, Republic of	
	KU Kuwait	
	LK Sri Lanka	
	LU Luxembourg	
	LV Latvia	
	LY Libya	
	MA Morocco	
	MC Monaco	
	MD Moldova	
	ME Montenegro	
	MG Madagascar	
	MH Marshall Islands	
	MI Micronesia	
	ML Mali	
	MM Myanmar	
	MN Mongolia	
	MO Macao	
	MP Northern Mariana Islands	
	MQ Martinique	
	MR Mauritania	
	MS Montserrat	
	MU Mauritius	
	MV Maldives	
	MW Malawi	
	MX Mexico	
	MY Malaysia	
	MZ Mozambique	
	NA Namibia	
	NC New Caledonia	
	NE Niger	
	NG Nigeria	
	NL Netherlands	
	NI Nicaragua	
	NO Norway	
	NP Nepal	
	OA Oman	
	OC Oceania	
	OD Oldenburg	
	OE Outermost Regions of France	
	OF Overseas Countries and Territories of France	
	OG Ogaden	
	OH Ohio	
	OI Omani	
	OK Oklahoma	
	OL Olbia	
	OM Oman	
	ON Ontario	
	OO Other	
	OP Overseas Possessions of the United Kingdom	
	OR Oregon	
	OS Ossetian	
	OT Overseas Territory of the United Kingdom	
	OU Ouagadougou	
	OV Overseas	
	PB Pacific Basin	
	PC Panama	
	PD Palestinian	
	PE Peru	
	PF French Polynesia	
	PG Papua New Guinea	
	PH Philippines	
	PK Pakistan	
	PL Poland	
	PM Puerto Rico	
	PN Pitcairn	
	PO Portugal	
	PP Palau	
	PQ Paraguay	
	PR Puerto Rico	
	PS Palestine	
	PT Portugal	
	PY Paraguay	
	RH Rhode Island	
	RI Rhode Island	
	RJ Rio de Janeiro	
	RK Republic of Korea	
	RL Rumania	
	RM Romania	
	RS Russia	
	RU Russian Federation	
	SF South Florida	
	SG Singapore	
	SH Saint Helena	
	SI Slovenia	
	SJ Svalbard	
	SK Slovakia	
	SL Sierra Leone	
	SM San Marino	
	SN Senegal	
	SO Somalia	
	SP Spanish Sahara	
	SR Suriname	
	SS South Sudan	
	ST Sao Tome and Principe	
	SV El Salvador	
	SW Sweden	
	SY Syria	
	TB Tennessee	
	TC Turks and Caicos Islands	
	TD Chad	
	TE Texas	
	TF Tokelau	
	TH Thailand	
	TJ Tajikistan	
	TK Tokelau	
	TL Timor-Leste	
	TM Turkmenistan	
	TN Tunisia	
	TO Tonga	
	TP Taiwan	
	TR Turkey	
	TS Tristan da Cunha	
	TU Tuvalu	
	TV Tuvalu	
	TX Texas	
	TY Tyne and Wear	
	UA Ukraine	
	UB Utah	
	UC Uganda	
	UD Upper District	
	UE Uruguay	
	UF Upper Florida	
	UG Uganda	
	UH Hawaii	
	UI Upper Iowa	
	UK United Kingdom	
	UL Upper Louisiana	
	UM United States Minor Outlying Islands	
	UN United Nations	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CZ Czechia	LK Sri Lanka	TN Tunisia
DE Germany, Federal Republic of	LU Luxembourg	ZM Zambia
DK Denmark	MC Monaco	US United States of America
FI Finland	MG Madagascar	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CA Canada	CZ Czech Republic	
CH Switzerland	DE Germany, Federal Republic of	
CL Chile	DK Denmark	
CO Colombia	EE Estonia	
DE Germany, Federal Republic of	FI Finland	
DK Denmark	FR France	
FI Finland	GB United Kingdom	
	GR Greece	
	GU Guadeloupe	
	HK Hong Kong	
	IL Israel	
	IN India	
	IR Iran	
	IS Iceland	
	IT Italy	
	JP Japan	
	KE Kenya	
	KR Korea, Republic of	
	KU Kuwait	
	LK Sri Lanka	
	LU Luxembourg	
	LV Latvia	
	LY Libya	
	MC Monaco	
	MD Moldova	
	ME Montenegro	
	MG Madagascar	
	MH Marshall Islands	
	MI Micronesia	
	ML Mali	
	MM Myanmar	
	MN Mongolia	
	MO Macao	
	MP Northern Mariana Islands	
	MQ Martinique	
	MR Mauritania	
	MS Montserrat	
	MU Mauritius	
	MV Maldives	
	MW Malawi	
	MX Mexico	
	MY Malaysia	
	MZ Mozambique	
	NA Namibia	
	NC New Caledonia	
	NE Nigeria	
	NF Norfolk Island	
	NI Nicaragua	
	NL Netherlands	
	NO Norway	
	NP Nepal	
	OZ New Zealand	
	PB Palau	
	PE Peru	
	PF French Polynesia	
	PG Papua New Guinea	
	PH Philippines	
	PK Pakistan	
	PL Poland	
	PM Saint Pierre and Miquelon	
	PN Pitcairn Islands	
	PR Puerto Rico	
	PT Portugal	
	PY Paraguay	
	QA Qatar	
	RD Rwanda	
	RE Reunion	
	RU Russian Federation	
	SA Saudi Arabia	
	SC Seychelles	
	SE Sweden	
	SG Singapore	
	SH St. Helena	
	SI Slovenia	
	SJ Svalbard and Jan Mayen	
	SK Slovakia	
	SL Sierra Leone	
	SM San Marino	
	SN Senegal	
	SO Somalia	
	SR Suriname	
	SS South Sudan	
	ST Sao Tome and Principe	
	SV El Salvador	
	TD Chad	
	TF French Southern Territories	
	TG Togo	
	TH Thailand	
	TJ Tajikistan	
	TK Tokelau	
	TL Timor-Leste	
	TM Turkmenistan	
	TN Tunisia	
	TO Tonga	
	TR Turkey	
	TT Trinidad and Tobago	
	TV Tuvalu	
	UA Ukraine	
	UG Uganda	
	US United States of America	
	UY Uruguay	
	UZ Uzbekistan	
	VN Vietnam	
	VG Virgin Islands, British	
	VI Virgin Islands, U.S.	
	VN Vietnam	
	WF Wallis and Futuna	
	YE Yemen	
	ZA South Africa	
	ZM Zambia	
	ZW Zimbabwe	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CA Canada	CZ Czech Republic	
CH Switzerland	DE Germany, Federal Republic of	
CL Chile	DK Denmark	
CO Colombia	EE Estonia	
DE Germany, Federal Republic of	FI Finland	
DK Denmark	FR France	
FI Finland	GB United Kingdom	
	GR Greece	
	GU Guernsey	
	HK Hong Kong	
	IL Israel	
	IN India	
	IR Iran	
	IS Iceland	
	IT Italy	
	JP Japan	
	KE Kenya	
	KR Korea, Republic of	
	KU Kuwait	
	LX Luxembourg	
	MC Monaco	
	MG Madagascar	
	MT Malta	
	MU Mauritius	
	NE Nigeria	
	NI Nicaragua	
	NL Netherlands	
	NO Norway	
	OZ New Zealand	
	PB Pakistan	
	PE Peru	
	PG Papua New Guinea	
	PH Philippines	
	PK Pakistan	
	PL Poland	
	PT Portugal	
	QA Qatar	
	RE Réunion	
	RU Russian Federation	
	SA Saudi Arabia	
	SC Seychelles	
	SE Sweden	
	SG Singapore	
	SI Slovenia	
	SJ Svalbard and Jan Mayen	
	SK Slovakia	
	SN Senegal	
	SO Somalia	
	SR Suriname	
	SS South Sudan	
	ST São Tomé and Príncipe	
	SV El Salvador	
	TD Chad	
	TH Thailand	
	TJ Tajikistan	
	TM Turkmenistan	
	TN Tunisia	
	TR Turkey	
	TT Trinidad and Tobago	
	TV Tuvalu	
	UA Ukraine	
	UG Uganda	
	US United States of America	
	UY Uruguay	
	UZ Uzbekistan	
	VC Saint Vincent and the Grenadines	
	VE Venezuela	
	VG Virgin Islands, British	
	VI Virgin Islands, U.S.	
	VN Vietnam	
	WF Wallis and Futuna	
	YE Yemen	
	ZA South Africa	
	ZM Zambia	
	ZW Zimbabwe	

[illegible]

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CA Canada	LA Laos	
CH Switzerland	LV Latvia	
CN China	LU Luxembourg	
DE Germany, Federal Republic of	MC Monaco	
DK Denmark	MG Madagascar	
FI Finland		

- 1 -

PROTEASE INHIBITORS

Background of the Invention

This invention relates to protease inhibitors, and particularly to transition state analogs.

5 Transition state analogs, compounds which are thought to resemble the substrates of enzymes, are thought to bind more tightly to the enzymes than the substrates themselves. Transition state analogs form complexes with enzymes at their catalytic sites.


10 Baugh et al. (Proteinases and Tumor Invasion ed. Strauli et al., Raven Press, New York, 1980, p./165) state that transition state analogs containing boronic acid moieties or aldehydes form tetrahedral adducts with serine proteases and are thus good inhibitors of these
15 enzymes. Further, they state that some peptide aldehydes have been synthetically prepared, that most are of microbial origin, and that "it would appear that changing the R-group [of synthetic peptides] to satisfy the specific requirements of a given protease should
20 result in both potent and specific inhibitors." They also state that transition state analogs containing cyclic ester moieties have been used to inhibit chymotrypsin and that "variations thereof may become useful as inhibitors of cathepsin G."

25 Yoshimoto et al. (J. Biochem .98: 975, 1985) describe prolyl endopeptidase inhibitors containing a protinal moiety. These inhibitors appear to act non-competitively.

Shenvi et al., U.S. Patent No. 4,499,082,
30 describe peptides having an α -amino boronic acid

- 2 -

residue. These peptides are reversible inhibitors of elastase. They have the structural formula




where R_2 is an alkyl group of one to six carbons which may have an aromatic substituent or an in-chain bivalent group.


Summary of the Invention

In a first aspect, the invention features compounds having the structure

10



and salts thereof, where T is a boronate group of the formula



where each D^1 and D^2 , independently, is a hydroxyl group or a group which is capable of being hydrolyzed to a hydroxyl group in aqueous solution at physiological pH; or T is a group of the formula



where G is either H, F or an alkyl group containing 1 to about 20 carbon atoms and optional heteroatoms which can be N, S, or O; or T is a phosphonate group of the formula




- 3 -

alkyl (each containing about 1-20 carbon atoms) and, optionally, heteroatoms which can be N, S, or O; where T is a group able to form a complex with the catalytic site of an enzyme); X includes one or more amino acids,


5 Y is



or



or



and each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and
 10 R^8 , independently, is a group which does not interfere significantly (i.e., does not raise the K_i of the compound to greater than 10^{-5}M) with site-specific recognition of the compound by the enzyme, while permitting a complex to be formed between the compound
 15 and the enzyme.

In preferred embodiments, T is a boronate group, each D^1 and D^2 , independently, is OH or F or D^1 and D^2 together form a ring containing 1 to about 20 carbon atoms, and optionally heteroatoms which can be
 20 N, S, or O; each R^{1-8} is H; X mimics the substrate recognised by the enzyme, for example X is pro-, thr-pro-, ala-pro-, ala-ala-pro-, ser-thr-pro-, pro-ser-, pro-thr-, or ser-pro- (pro = proline, thr = threonine, and ser = serine); X contains both an amino
 25 acid and a blocking group, such as an acetyl group; the enzymes inhibited by the compounds of the invention are most preferably serine

- 4 -

proteases, even more preferably IgA1 proteases; and the analog has a binding constant of at least $10^{-7}M$, most preferably $10^{-10}M$.

In a second aspect, the invention features a
5 compound, which is useful as an intermediate in the synthesis of compounds of Formula (1), having the formula




where V is $(\text{CH}_3)_3\text{Si}-$ or $\text{H}-$, Y is



10



or



each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8
is independently H, or C_1 - C_{10} alkyl or aryl, and Z is



15 where D^1 and D^2 are as defined above.

The invention also features a method for producing the above compounds. The method includes the steps of reacting a compound of the formula

- 5 -

where Q is Cl or Br, and R², Y, Z, and D¹ and D²
are as defined above, with a compound of the formula



in an inert solvent at a temperature between -78°C and
5 25°C, and heating the reaction mixture to at least 80°C.

In preferred embodiments the reacting step is
performed in the absence of an inert solvent.

The compounds of the invention are peptide
derivatives (or intermediates in their formation) which
10 are potent inhibitors of proteolytic enzymes (especially
those produced by pathogens) which are post-prolyl
cleaving enzymes, e.g., the serine proteases able to act
on IgA1 proteins. The compounds generally have a
proline, proline analog, a 2-azetidinecarboxylic acid,
15 or pipecolic acid linked to a group, for example a
boronate group, which mimics the transition state of an
enzyme substrate, and to a peptide moiety which
preferably mimics the site of the substrate acted upon
by the enzyme of interest. It is proposed that the
20 peptide moiety is recognized as a substrate by the
enzyme to be inhibited, and it then enters an active
site, catalytic site, or transition state binding site
of the enzyme, and the transition state-mimicing group
of the compound of the invention binds strongly at this
25 site. This binding advantageously prevents the enzyme
from acting upon its natural substrate. The high
affinity of these compounds make them effective at
concentrations as low as 10⁻⁷M, or even 10⁻¹⁰M.

Thus, the present invention also provides
30 compositions including one or more compounds of formula
(1) above, and methods of using such compounds or
compositions in the inhibition of post-prolyl cleaving

- 6 -

Other features and advantages of the invention will be apparent from the following description of preferred embodiments thereof, and from the claims.

Description of the Preferred Embodiments

5 The Figures will first briefly be described.

Drawing

Figure 1 shows the general formula of three preferred compounds.

Figure 2 is a diagrammatic representation of
10 the synthesis of a boroProline compound.

Structure

The compounds of the invention have the general structure recited in the Summary of the Invention above. Examples of preferred structures are those
15 referred to as preferred embodiments above.

The structure of the compounds is such that at least a portion of the amino acid sequence near the cleavage site of a post-prolyl cleaving enzyme substrate is duplicated, or nearly duplicated. This duplication
20 is in part responsible for the ability of the compounds to inhibit the enzyme, by a mechanism thought to involve competitive inhibition between the compound and the actual enzyme substrate.

The choice of amino acid sequence affects the
25 inhibitory activity of the compound, and its specificity. As a first step in determining a suitable sequence, the amino acid sequence of the substrate is determined near its cleavage site. In some cases, such as for serine proteases, a suitable inhibitor sequence
30 is the amino acid sequence N-terminal to (i.e., to the left of) the proline cleavage point, and includes that proline. Peptide fragments can be synthesized and then tested to determine their efficacy as inhibitors, using

- 7 -


similar fashion, by testing the inhibitory effect of a particular compound on a variety of enzyme activities. The compounds preferably inhibit the activity of enzymes detrimental to an animal or human patient, and

5 preferably do not inhibit necessary enzymes.


The compounds of the invention can be designed so that they are resistant to attack by agents which might otherwise cause their catabolic degradation by cleavage of one or more peptide bonds in the peptides.

10 For example, an N-terminal blocking group, such as acetyl, can increase the half-life of the compounds in vivo, and thus improves inhibition.

The structure of such blocking groups can vary widely. In one blocking reaction, a hydrogen atom of
15 the amino terminal amino group is replaced, generally in a dehydration reaction. Thus, blocking groups such as



readily added to a peptide chain. Others include



20 Such N-terminal blocking groups may be employed not only to protect amino-terminal groups, but also may protect-side chain amino group of amino acids which make up X, such as where X includes Lys or Arg. Similarly, amino acid residues having acidic or hydroxy side chains
25 can be protected in the form of t-butyl, benzyl, or other suitable esters or ethers. Short length of the compounds of the invention (2-7, preferably 3 or 4 amino acids) is advantageous because it provides stability and

- 8 -

The compounds also include a group (T) which causes the compound to complex with an enzyme, not only in a competitive fashion, but in a chemically reactive manner to form a strong bond between the compound and the enzyme. This group thus acts to bind the compound to the enzyme, and increases the inhibitory binding constant (K_i) of the compound. Examples of such groups include boronates, fluoroalkyl ketones and substituted phosphonates (of the formulae given in the Summary above, examples of which are shown in Fig. 1). These groups are covalently bonded to the prolyl residue of the compound, as in the above formula.

The proline or proline analog, represented by



above, is chosen so that it mimics the structure of proline recognized by the active site of the enzyme to be inhibited. It can be modified by providing R^{1-2} groups which do not interfere significantly with this recognition and thus do not significantly affect the K_i of a compound. Thus, one or more hydroxyl groups can be substituted to form hydroxy-proline, and methyl or sugar moieties may be linked to these groups. One skilled in the art will recognize that these groups are not critical in this invention and that a large choice of substituents are acceptable for R.

Examples of compounds having utility as serine protease inhibitors include compounds having a peptide chain similar to the substrate of IgA1 and IgA2 proteases. Plaut 1983 Ann Rev. Microbiol. 37,

- 9 -

bonds of IGA, the immunoglobulin that provides antibody defense of mucosal surfaces, resulting in a nonfunctional immunoglobulin and impairment of the host defense system. This is expected to be a strong
5 contributing factor to pathogenesis of organisms such as Streptococcus sanguis, S. pneumoniae, Neisseria gonorrhoeae, N. meningitidis, and Haemophilus influenzae.

IgA1 proteases recognize the cleavage site Ser-Thr-Pro-Pro-X (where X is any amino acid),
10 hydrolyzing between Pro and X (i.e., they are post-prolyl cleaving enzymes). Accordingly, Ser-Thr-Pro-Pro-T is a suitable compound of the invention for inhibiting IGA1 proteases. The Ser or Thr in this compound can be readily substituted with any of
15 the 20 naturally occurring amino acids, most preferably those having non-bulky side groups, such as Ala and Gly. It is also possible to substitute non-naturally occurring amino acids, such as 2-azetidinecarboxylic acid or pipecolic acid (which have six-membered, and
20 four-membered ring structures respectively) for either of the Pro residues. Those skilled in the art will recognize that there are other such changes which can be made without significantly affecting the inhibitory character of these compounds.

25 In the case of type 1 IgA proteases, the cleavage site is Pro-Thr-Pro-X, with hydrolysis occurring between Pro and X. Thus, a preferred compound of the invention for inhibiting IGA2 proteases has the formula Pro-Thr-Pro-T. Thr can be substituted by any of
30 the naturally occurring amino acids, especially ones having non-bulky side groups, such as Ala, Gly, or Ser.

Other examples of enzymes which can be inhibited according to the invention include other post

- 10 -

degrading enzymes, vasopressin degrading enzymes, and oxytocin degrading enzymes. Further, the serine protease, dipeptidyl peptidase Type IV, on T-lymphocytes (Andrews et al., Clin. Lab. Haemato. 7: 359-368, 1985), which plays a role in regulation of the immune response, can be inhibited by suitable such compounds. These inhibitors may be useful in treatment of AIDS. Walters et al., Molecular and Cellular Biochemistry 30: 111-127, 1980 describes other such enzymes, and is hereby incorporated by reference.

Synthesis

Synthesis of boroProline

Referring to Fig. 2, the starting compound I is prepared essentially by the procedure of Matteson et al., 3 Organometallics 1284, 1984, except that a pinacol ester is substituted for the pinanediol ester. Similar compounds such as boropiecolic acid and 2-azetidine boronic acid can be prepared by making the appropriate selection of starting material to yield the pentyl and propyl analogs of compound I. Further, Cl can be substituted for Br in the formula, and other diol protecting groups can be substituted for pinacol in the formula, e.g., 2,3-butanediol and alpha-pinanediol.

Compound II is prepared by reacting compound I with $[(CH_3)_3O_3Si]_2N^-Li^+$. In this reaction hexamethyldisilazane is dissolved in tetrahydrofuran and an equivalent of n-butyllithium added at $-78^\circ C$. After warming to room temperature ($20^\circ C$) and cooling to $-78^\circ C$ an equivalent of compound I is added in tetrahydrofuran. The mixture is allowed to slowly come to room temperature and to stir overnight. The alpha-bis[trimethylsilane]-protected amine is isolated by evaporating solvent and adding hexane under anhydrous

- 11 -

under a nitrogen blanket, yielding a hexane solution of compound 2.

Compound III, the N-trimethylsilyl protected form of boroProline is obtained by the thermal
5 cyclization of compound II during the distillation process in which compound II is heated to 100-150°C and distillate is collected which boils 66-62°C at 0.06-0.10 mm pressure.

Compound IV, boroProline-pinacol hydrogen
10 chloride, is obtained by treatment of compound III with HCl:dioxane. Excess HCl and by-products are removed by trituration with ether. The final product is obtained in a high degree of purity by recrystallization from ethyl acetate. H-boroProline as the hydrochloride salt
15 is preferred, but other salts forms such as the hydrobromide and trifluoroacetate can readily be obtained by substitution of the appropriate acid for HCl.

The boroProline esters can also be obtained by treatment of the reaction mixture obtained in the
20 preparation of compound II with anhydrous acid to yield 1-amino-4-bromobutyl boronate pinacol as a salt. Cyclization occurs after neutralizing the salt with base and heating the reaction.

Synthesis of boroProline Peptides

25 General methods of coupling of N-protected peptides and amino acids with suitable side-chain protecting groups to H-boroProline-pinacol are applicable. When needed, side-chain protecting and N-terminal protecting groups can be removed by treatment
30 with anhydrous HCl, HBr, trifluoroacetic acid, or by catalytic hydrogenation. These procedures are known to those skilled in the art of peptide synthesis. One exception is that in the preparation of a compound with

- 12 -

protecting groups from threonine hydroxyl group results in a complex mixture of products. Thus, the use of hydrogenolytic protecting groups for threonyl residue is preferred.

5 The mixed anhydride procedure of Anderson et al., J. Am. Chem. Soc., 89:5012 (1984) is preferred for peptide coupling. The mixed anhydride of an N-protected amino acid or a peptide varying in length from a dipeptide to tetrapeptide is prepared by dissolving the
10 peptide in tetrahydrofuran and adding one equivalent of N-methylmorpholine. The solution is cooled to -20°C and an equivalent of isobutyl chloroformate is added. After 5 minutes, this mixture and one equivalent of triethylamine (or other sterically hindered base) are
15 added to a solution of H-boroPro-pinacol dissolved in either cold chloroform or tetrahydrofuran.

 The reaction mixture is routinely stirred for one hour at -20°C and 1-2 hours at room temperature (20°C). Solvent is removed by evaporation, and the
20 residue is dissolved in ethyl acetate. The organic solution is washed with 0.20 N hydrochloric acid, 5% aqueous sodium bicarbonate, and saturated aqueous sodium chloride. The organic phase is dried over anhydrous sodium sulfate, filtered, and evaporated. Products are
25 purified by either silica gel chromatography or gel permeation chromatography using Sephadex™ LH-20 and methanol as a solvent.

 Previous studies have shown that the pinacol protecting group can be removed in situ by preincubation
30 in phosphate buffer prior to running biological experiments; Kettner et al., J. Biol. Chem. 259: 15106-15114 (1984). Several other methods are also applicable for removing pinacol groups from peptides

- 13 -

product. First, the peptide can be treated with diethanolamine to yield the corresponding diethanolamine boronic acid ester, which can be readily hydrolyzed by treatment with aqueous acid or a sulfonic acid substituted polystyrene resin as described in Kettner et al., Id. Both pinacol and pinanediol protecting groups can be removed by treating with BCl_3 in methylene chloride as described by Kinder et al., J. Med. Chem., 28: 1917. Finally, the free boronic acid can be converted to the difluoroboron derivative ($-\text{BF}_2$) by treatment with aqueous HF as described by Kinder et al., Id.

Similarly, different ester groups can be introduced by reacting the free boronic acid with various di-hydroxy compounds (for example, those containing heteroatoms such as S or N) in an inert solvent.

The following abbreviations are used in the examples below. THF - tetrahydrofuran; H-Pro-OBzl - the benzyl ester of proline; H-Thr (OBzl)-OH - the benzyl ether derivative of threonine; Boc - the tertiary butyloxycarbonyl group; FABMS - fast atom bombardment mass spectrometry.

All natural amino acids are in the L-configuration. H-boroProline is in the D,L-configuration.

Example 1: Preparation of boroProline-pinacol
(H-boroPro-pinacol)

The intermediate, 4-Bromo-1-chlorobutyl boronate pinacol, was prepared by the method in Matteson et al., Organometallics, (3): 1284-1288 (1984), except that conditions were modified for large scale preparations and the pinacol was substituted for the pinanediol protecting group.

- 14 -

3-bromopropyl boronate pinacol was prepared by hydrogenboration of allyl bromide (173 ml, 2.00 moles) with catechol borane (240 ml, 2.00 moles). Catechol borane was added to allyl bromide and the reaction
5 heated for 4 hours at 100°C under a nitrogen atmosphere. The product, 3-bromopropyl boronate catechol (bp 95-102°C, 0.25 mm), was isolated in a yield of 49% by distillation. The catechol ester (124 g, 0.52 moles) was transesterified with pinacol (61.5 g, 0.52
10 moles) by mixing the component in 50 ml of THF and allowing them to stir for 0.5 hours at 0°C and 0.5 hours at room temperature. Solvent was removed by evaporation and 250 ml of hexane added. Catechol was removed as a crystalline solid. Quantitative removal was achieved by
15 successive dilution to 500 ml and to 1000 ml with hexane and removing crystals at each dilution. Hexane was evaporated and the product distilled to yield 177 g (bp 60-64°C, 0.35 mm).

4-Bromo-1-chlorobutyl boronate pinacol was
20 prepared by homologation of the corresponding propyl boronate. Methylene chloride (50.54 ml, 0.713 moles) was dissolved in 500 ml of THF, 1.54 N n-butyllithium in hexane (480 ml, 0.780 moles) was slowly added at -100°C. 3-Bromopropyl boronate pinacol (178 g, 0.713
25 moles) was dissolved in 500 ml of THF, cooled to the freezing point of the solution, and added to the reaction mixture. Zinc chloride (54.4 g, 0.392 moles) was dissolved in 250 ml of THF, cooled to 0°C, and added to the reaction mixture in several portions. The
30 reaction was allowed to slowly warm to room temperature and to stir overnight. Solvent was evaporated and the residue dissolved in hexane (1 liter) and washed with water (1 liter). Insoluble material was discarded.

- 15 -

filtering, solvent was evaporated. The product was distilled to yield 147 g (bp 110-112°C, 0.200 mm).

N-Trimethylsilyl-boroProline pinacol was prepared first by dissolving hexamethyldisilazane (20.0 g, 80.0 mmoles) in 30 ml of THF, cooling the solution to -78°C, and adding 1.62 N n-butyllithium in hexane (49.4 ml, 80.0 mmoles). The solution was allowed to slowly warm to room temperature. It was re-cooled to -78°C and 4-bromo-1-chlorobutyl boronate pinacol (23.9 g, 80.0 mmoles) added in 20 ml of THF. The mixture was allowed to slowly warm to room temperature and to stir overnight. Solvent was removed by evaporation and dry hexane (400 ml) added to yield a precipitant which was removed by filtration under an nitrogen atmosphere. The filtrate was evaporated and the residue distilled, yielding 19.4 g of the desired product (bp 60-62°C, 0.1-0.06 mm).

H-boroProline-pinacol.HCl was prepared by cooling N-trimethylsilyl-boroProline-pinacol (16.0 g, 61.7 mmoles) to -78°C and adding 4 N HCL:dioxane 46 ml, 185 mmoles). The mixture was stirred 30 minutes at -78°C and 1 hour at room temperature. Solvent was evaporated and the residue triturated with ether to yield a solid. The crude product was dissolved in chloroform and insoluble material removed by filtration. The solution was evaporated and the product crystallized from ethyl acetate to yield 11.1 g of the desired product (mp 156.5-157°C).

Example 2: Preparation of Boc-Ala-Pro-boroPro-pinacol

Boc-Ala-Pro-boroPro-pinacol was prepared by coupling Boc-Ala-Pro-OH to H-boroPro-pinacol. First, the dipeptide, Boc-Ala-Pro-OBzl, was prepared by the mixed anhydride procedure. Boc-Ala-OH (10g, 52.8 mmoles) was reacted with N-methylmorpholine (5.8 ml

- 16 -

52.8 mmole) and isobutyl chloroformate (6.8 ml, 52.8 mmole) for 5 minutes in 50 ml of THF at -20° . The reaction mixture and additional N-methylmorpholine (5.8 ml) were added to H-Pro-OBzl.HCl (12.8 g, 52.8 mmole) dissolved in 50 ml of cold chloroform. After the mixture was stirred for 1 hour at -20°C and 2 hours at room temperature, it was filtered and the filtrate evaporated. The residue was dissolved in ethyl acetate and washed sequentially with 0.2 N hydrochloric acid, 5% aqueous NaCO_3 , and saturated aqueous NaCl. The organic phase was dried over anhydrous Na_2SO_4 , filtered, and evaporated to yield Boc-Ala-Pro-OBzl as an oil (14.1 g). The benzyl ester protecting group was removed by dissolving Boc-Ala-Pro-OBzl (35 g, 92.8 mmole) in 200 ml of methanol and hydrogenating for 1 hour in the presence of 0.5 g of 10% Pd/C. The catalyst was removed by filtration and the solvent evaporated. The residue was crystallized from ethyl acetate to yield 16.1 g of Boc-Ala-Pro-OH (mp $153.5\text{--}154.5^{\circ}\text{C}$).

Boc-Ala-Pro-OH (1.26g, 4.28 mmole) was coupled to H-boroPro-pinacol by the general procedure described for the preparation of Boc-Ala-Pro-OBzl. Boc-Ala-Pro-OH (1.26 g, 4.28 mmole) was dissolved in 10 ml of THF and cooled to -20°C ; N-methylmorpholine (0.47 ml, 4.28 mmole) and isobutyl chloroformate (0.557 ml, 4.28 mmole) were added. After stirring for 5 minutes, 10 ml of cold THF and triethylamine (0.597 ml, 4.28 mmole) were added and the mixture added to a cold solution of H-boroPro-pinacol.HCl (1.0 g, 4.28 mmole) in 5 ml chloroform. After dissolving the reaction product in ethyl acetate and washing with aqueous HCl, NaHCO_3 , and saturated aqueous NaCl, 0.39 g of material were obtained. It was further purified by chromatography on

- 17 -

Example 3: Preparation of H-Ala-Pro-
boroPro-pinacol.HCL

Boc-Ala-Pro-boroPro-pinacol (0.58 g, 1.25 mmoles) was allowed to react with 2.5 mol of 4 N HCl:dioxane for 30 minutes at room temperature. Ether (50 ml) was added to yield 0.22 g of amorphous white solid.

Example 4: Preparation of Ac-Ala-Pro-boroPro-pinacol

H-Ala-Pro-boroPro-pinacol.HCL (0.22 g, 0.55 mmoles) was dissolved in 3 ml of THF and cooled to 0°C. Acetic anhydride (0.078 ml, 0.825 mmoles) and triethylamine (0.115 ml, 0.825 mmoles) were added and the reaction was allowed to come to room temperature. After approximately 25 minutes, additional triethylamine (0.25 ml, 0.179 mmoles) was added. After a total reaction time of 45 minutes, the reaction solution was applied to a 2.5 X 100 column of LH-20 in methanol and fractions (approximately 7 ml) collected. Fraction 22-28 contained 0.16 g of the desired product.

Example 5: Preparation of MeOSuc-Ala-
Ala-Pro-boroPro-pinacol.

MeOSuc-Ala-Ala-Pro-OH was prepared by the procedure described in Kettner et al., J. Biol. Chem., 259: 15106-15114 (1984). MeOSuc-Ala-Ala-Pro-OH (1.59 g, 4.28 mmoles) was coupled to H-boroPro-pinacol.HCl (1.00 g, 4.28 mmoles) by the mixed anhydride procedure described for the preparation of Boc-Ala-Pro-boroPro-pinacol except that, after filtration and evaporation of the reaction solvent, it was applied to a 2 cm column containing 10 g of silica gel equilibrated with chloroform. The column was eluted with chloroform and fractions containing the desired product were evaporated and triturated with hexane to yield 0.58 g of a white solid.

- 18 -

Example 6: Preparation of Boc-Pro-
Thr(OBzl)-boroPro-pinacol.

- Boc-Pro-Thr(OBzl)-OH was prepared by coupling Boc-Pro-OSu (the N-hydroxysuccinamide ester of
- 5 Boc-Pro-OH) to H-Thr (OBzl) - OH. H-Thr (OBzl)-OH.HCL was dissolved in 25 ml of water, NaHCO_3 (6.21 g, 73.9 mmoles) and a solution of Boc-Pro-OSu (5.07 g, 16.25 mmoles) in 25 ml of dioxane were added. After stirring 3 hours, the reaction mixture was acidified with
- 10 hydrochloric acid and the product extracted into ethyl acetate. It was washed with 0.2 N HCl prepared in saturated aqueous NaCl, saturated NaCl and dried over anhydrous Na_2SO_4 . Solvent was evaporated to yield 6.24 g (mp 120-12.5°).
- 15 Boc-Pro-Thr(OBzl)-boroPro-pinacol was prepared by coupling Boc-Pro-Thr(OBzl)-OH (2.70 g, 6.42 mmoles) to H-boroPro-pinacol.HCL (1.50 g, 6.42 mmoles) using the procedure described for Boc-Ala-Pro-boroPro-pinacol. The product (2.4 g) was purified by chromatography on a
- 20 2.5 X 50 cm column of LH-20 in methanol and was obtained as 0.84 g of oil.

Example 7: Preparation of Boc-Pro-Thr-boroPro-Pinacol

- Boc-Pro-Thr-boroPro-pinacol was prepared by hydrogenation of Boc-Pro-Thr(OBzl)-boroPro-pinacol (from
- 25 Example 6, 0.585 g, 0.79 mmoles). The protected peptide was dissolved in 100 ml of methanol and was hydrogenated for 2 hours on a Parr apparatus in the presence of 0.5 g of 10% Pd/C. The reaction was filtered and solvent evaporated. The product was allowed to stir with
- 30 hexane, and hexane insoluble material removed by filtration. The filtrate was evaporated to yield the desired product (0.24 g) as a white foam.

Use

- 19 -

administered in an effective amount either alone or in combination with a pharmaceutically acceptable carrier or diluent.

The compounds of formula 1 or compositions
5 thereof can be used to treat mammals, e.g., humans, suffering from or subject to infections by pathogenic bacteria which produce deleterious post prolyl cleaving enzymes, e.g., the IgA1 protease of Haemophilus influenza. The compounds or compositions can be
10 administered alone or in combination with one another, or in combination with other therapeutic agents. The dosage level may be 1-500 mg/kg/day of the formula 1 compounds.


When administered to mammals (e.g., orally,
15 topically, intramuscularly, intraperitoneally, intravenously, parenterally, nasally or by suppository), the compounds of the invention enhance the ability of e.g., the immune system of the mammal to fight bacteria which produce an IgA-1 protease, thus slowing the course
20 of infection by the bacteria.

Other embodiments are within the following claims.

- 20 -

Claims

1. A compound, having the structure




wherein T is a group of the formula



where each D^1 and D^2 , independently, is a hydroxyl group or a group which is capable of being hydrolysed to a hydroxyl group in aqueous solution at physiological pH; a group of the formula



where G is either H, F or an alkyl group containing 1 to about 20 carbon atoms and optional heteroatoms which can be N, S, or O; or a phosphonate group of the formula






15 where each J, independently, is O-alkyl, N-alkyl, or alkyl, each said O-alkyl, N-alkyl or alkyl comprising about 1-20 carbon atoms and, optionally, heteroatoms which can be N, S, or O; said T being able to form a complex with the catalytic site of a post-prolyl

20 cleaving enzyme;

X comprises an amino acid,

- 21 -

Y is



- 5 and each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 , separately is a group which does not significantly interfere with site specific recognition of said compound by said enzyme, and allows said complex to be formed with said enzyme.
- 10 2. The compound of claim 1, wherein T is a boronate group.
3. The compound of claim 1, wherein T is a substituted phosphonate group or a fluoroalkyl ketone group.
- 15 4. The compound of claim 1 wherein each R^{1-8} is H.
5. The compound of claim 1 wherein X mimics the site of a substrate recognized by said enzyme.
6. The compound of claim 1 wherein X comprises
- 20 a blocking group.
7. The compound of claim 6 wherein X comprises an acetyl group.
8. The compound of claim 5 where X is pro-, thr-pro-, ala-pro-, ala-ala-pro-, ser-thr-pro-,
- 25 pro-ser-, pro-thr-, or ser-pro-.
9. The compound of claim 1, said enzyme being a serine protease.

- 22 -

10. The compound of claim 9, said enzyme being a serine protease.


11. The compound of claim 8, said enzyme being an IgA1 protease.

5 12. The compound of claim 1, wherein said compound has a binding constant to said enzyme of at least 10^{-10} M.


13. The compound of claim 10, wherein said compound has a binding constant of at least 10^{-7} M.

10 14. The compound of claim 1 admixed within a pharmaceutically acceptable carrier substance.


15. A compound having the formula



where V is $(\text{CH}_3)_3\text{Si}-$ or $\text{H}-$, Y is



or



each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 is independently H, or C_1 - C_{20} alkyl or aryl, and Z is




where each D^1 , D^2 independently is a group which is


- 23 -

16. The compound of claim 1 or 14 wherein, each D^1 and D^2 is, independently, F or D^1 and D^2 together are a ring containing 1 to about 20 carbon atoms, and optionally heteroatoms which can be N, S, 5 or O.

17. A method for producing the compound of claim 15, comprising the steps of reacting



where Q is Cl or Br, and R^2 is H or C_1-C_{10} alkyl 10 or aryl, with



in an inert solvent at a temperature between $-78^\circ C$ and $25^\circ C$, and

heating the reaction mixture to at least $80^\circ C$.


15 18. The method of claim 17 wherein said reacting is in the presence of an inert solvent.

19. A method for inhibiting a post prolyl cleaving enzyme in a mammal, comprising administering to said mammal an effective amount of a compound of claim 1.


20 20. The method of claim 19 wherein said amount is 1-500 mg/kg/day.

Abstract of the Disclosure


A compound having the structure



where T is of the formula $\begin{array}{c} \text{D}^2 \\ | \\ -\text{B}-\text{D}^1 \end{array}$, where each D^1 and
 5 D^2 , independently, is a hydroxyl group or a group
 which is capable of being hydrolysed to a hydroxyl group
 in aqueous solution at physiological pH; a group of the
 formula




10 where G is either H, F or an alkyl group containing 1 to
 about 20 carbon atoms and optional heteroatoms which can
 be N, S, or O; or a phosphonate group of the formula




where J is O-alkyl, N-alkyl, or alkyl, each comprising
 15 about 1-20 carbon atoms and, optionally, heteroatoms
 which can be N, S, or O; T being able to form a complex
 with the catalytic site of an enzyme, X is a group
 having at least one amino acid,


Y is



or




or




5 and each R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 is separately a group which does not interfere significantly (i.e., does not lower the K_i of the compound to less than $10^{-7}M$) with site-specific recognition of the compound by the enzyme, and allows a
10 complex to be formed with the enzyme.


1 / 2



Prolyl Boronate




Prolyl Trifluoro alkyl ketone




Prolyl phosphonate

FIG 1



2 / 2



4-bromo-1-chlorobutyl boronate pinacol



4-bromo-1-[(bistrimethylsilyl) amino] butyl boronate pinacol



INTERNATIONAL SEARCH REPORT

International Application No. PCT/US88/03506

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) * According to International Patent Classification (IPC) or to both National Classification and IPC INT. CL. 4th Ed. A61K 37/02; C07K 5/06, 5/08, 5/10; see attn: US. CL. 514/18, 19; 530/329, 330, 331; 548/517, 518, 533, 540						
II. FIELDS SEARCHED <div style="text-align: center;">Minimum Documentation Searched †</div> <table style="width: 100%;"> <tr> <td style="width: 30%;">Classification System ‡</td> <td style="width: 70%;">Classification Symbols</td> </tr> <tr> <td style="height: 40px; vertical-align: top;">US</td> <td style="vertical-align: top;">514/18, 19; 530/329, 330, 331; 548/517, 518, 533, 540</td> </tr> </table> <div style="text-align: center; font-size: small;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ‡</div>			Classification System ‡	Classification Symbols	US	514/18, 19; 530/329, 330, 331; 548/517, 518, 533, 540
Classification System ‡	Classification Symbols					
US	514/18, 19; 530/329, 330, 331; 548/517, 518, 533, 540					
III. DOCUMENTS CONSIDERED TO BE RELEVANT ††						
Category *	Citation of Document, †† with indication, where appropriate, of the relevant passages ††	Relevant to Claim No. ††				
A	US, A, 4,652,552 (Kettner) Published 24 March 1987	1-20				
A	US, A, 4,636,492 (Kettner) Published 13 January 1987	1-20				
A	US, A, 4,644,055 (Kettner) Published 17 February 1987	1-20				
A	US, A, 4,582,821 (Kettner) Published 15 April 1986	1-20				
A	US, A, 4,499,082 (Shenvi) Published 12 February 1985	1-20				
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: ††</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>						
IV. CERTIFICATION						
Date of the Actual Completion of the International Search ‡		Date of Mailing of this International Search Report ‡				
<div style="text-align: center;"> International Searching Authority </div>		<div style="text-align: center;"> AUTHORIZED OFFICER </div>				

PCT/US88/03506

CLASSIFICATION OF SUBJECT MATTER

INT. CL. 4th Ed. CO7D 405/00, 207/00

PCT/US88/03506

CLASSIFICATION OF SUBJECT MATTER

INT. CL. 4th Ed. CO7D 405/00, 207/00